

# The Effect of Health Literacy on Self-Management Related to Food Consumption Among Older Adults with Hypertension in The Community

Rungnapa Pongkiatchai\*, Sresuda Wongwisadkul & Orranuch Chusri

Department of community Nursing, Faculty of Nursing Suan Dusit University, Bangkok, 10300 Thailand

## Abstract

The study aimed to study the effect of the health literacy program on self - management related to food consumption among older adults with hypertension in the community. The study design was quasi-experimental study with pre and posttest groups. The participants consisted of 64 older adults with hypertension who were randomly assigned to an experimental and control group. Participants in experimental group received health education and six sessions of health literacy. Each session lasted 20-60 minutes within eight weeks. Control group received usual care and health education from health provider. self-administration questionnaire (nutrition literacy, food consumption pattern and self-management) was used to collect the data. Data were analyzed using independent t-test. Group reflection analyzed using content analysis. Results showed that mean score of nutritional literacy, food consumption, and self-management of the experiment group was significantly higher than the mean score of the control group (p-value <.001). Results from the reflection in the experimental group suggested some issues, including the understanding, access to information, and applying knowledge. Participants suggested the methods to enhance health literacy for self-management on food consumption including providing knowledge, communication, sharing experiences, and using photographic media. In sum, health literacy program can be used as a tool to enhance knowledge related to health.

**Keywords:** health literacy, self – management, older adults, hypertension, food consumption

## \*Corresponding Author

e-mail: [rungnapa\\_pon@dusit.ac.th](mailto:rungnapa_pon@dusit.ac.th)

## Introduction

The older adult population in Thailand is on the risk resulting in ageing society. It has been predicted that Thailand will become an aged society in 2030 (National Committee on the Older adult, Department of Older adults, Ministry of Social Development and Human Security, 2020). This will inevitably affect the shift of social services towards the older adult population as an important target group for health promotion. At an older age, changes occur both physically and mentally. The changes in the body reduce the ability to care for themselves and subsequently, they have to rely more on family care in terms of economic, social, and daily activities. However, the requirement for older adult care varies according to health conditions. Normal and at-risk older adult people can participate in outdoor or community activities due to less or no physical limitations. Moreover, the community can put their effort to participate in care for the older adult with complex illnesses. Care activities associated with health conditions, can be focused on health promotion and surveillance of risks to severity or complications of such diseases.

The significant health problems among the older adult include suffering from chronic diseases such as high blood pressure, diabetes, heart disease, etc. Chronic diseases are related to many modifiable risk factors such as unhealthy diets, physical inactivity, consumption of tobacco and alcohol, and being overweight or obese (Ekplakorn, (editor), 2021). Behaviour changes particularly in the retirement age group (from 60 onwards) prevent any further damage to their health. Moreover, any retirees without any health problem can also be benefitted from serious illnesses later on in life. In 2020, the Thai National Health Examination Survey (NHES VI) reported that the prevalence of older adult with hypertension was 60.7% and prevalence of hypertension in Bangkok metropolitan was 27.2%. Division of Non-communicable Diseases in Thailand also reported the number of deaths from uncontrolled hypertension at 9.303 per 100,000 population.

High blood pressure, or hypertension, is a major health problem in older adults. The vascular system changes with age resulting in stiffer arteries. This causes increasing blood pressure. Hypertension is known as "the silent killer," and often does not show any signs of illness. If hypertension cannot be controlled with lifestyle changes and medication, it can lead to serious health problems, including cardiovascular disease such as heart disease and stroke, vascular dementia, eye problems, and kidney disease (WHO, 2021). These complications had impact on older adult such as morbidity and mortality. These effects lead to dependency on family and community (Kanjanapibulwong et al., 2020; Ekplakorn, (editor), 2021). In addition, consumer behaviour can also affect health conditions in those who are considered well. It can cause changes in biomarkers to indicate a person is at risk of developing the disease. Therefore, blood pressure control needed the self-management. The World Health Organization (2013) recommended the involvement of patients through their own self-management surveillance such as smoking cessation, weight management, low-sodium and low-fat diet to better control high blood pressure. Self-management also applies to health promotion, and the distal outcomes are health status and quality of life. In addition, a systematic review found that the majority of hypertensive individuals with higher health literacy tend to have better blood pressure control (Mohd Isa et al., 2021)

Self-management is the intrinsically controlled ability of an active, responsible, informed and autonomous individual to live with the medical condition, role and emotional consequences of their chronic conditions in partnership with their social network and the healthcare providers (Van de Velde et al., 2019). Therefore, the adjustment of consumer behaviour is a good guideline for the self-management of the older adult. Meanwhile, advances in telecommunication technology have given the older adult access to information through various channels and a mass of health information. If older adults do not have sufficient health literacy, they could potentially struggle to comprehend an overflowing information and to make decision on healthy choice. The latter could lead to an inappropriate decision on healthy behaviours. On the other hand, the older adult with chronic illnesses manages their consumption appropriately with health literacy through various information and communication channels that will bring them a good quality of life according to their health condition. The development of health literacy and self-management is mutually complemented one another (Norris, et al., 2002; Kim., 2004; Chodosh, et al., 2005; Chao, et al., 2013; Wang, et al., 2017).

Health information education is a model of action to change consumer behaviour, and the outcome is the number of people who consumes that information. However, Nutbeam (2010) argues that the results of health information education should be the skill of individuals to make decisions or improve their behaviour. This is known "health literacy". Health literacy is considered as one of the most important skills to control people health. Health literacy is the set of patients' cognitive and social skills that allows them to access, understand, and use information in ways that promote and maintain their health (WHO, 1998). However, one specific form of health literacy is nutrition literacy. Nutrition literacy reflects the ability to access, interpret, and use nutrition information and exactly focuses on health literacy skills related to food consumption (Velardo, 2015). Food consumption is a periodic behavior. It is triggered at various moments of the day by a number of converging factors (time of day, need state, sensory stimulation, social context, etc.). As eating progresses, inhibitory influences of many origins (sensory, gastric, hormonal, neural, as well as cognitive) develop and finally bring the meal to an end (Bellisle, 2019). Previous research evaluated the relationship between health literacy and food consumption among adult population. The results showed that high food literacy was associated with increased consumption of fruits and vegetables (Namdar et al., 2021)

The educational program is blended with sharing and learning, and self-talk. It helps the older adult manage themselves to change dietary habits under their health condition. Developing health literacy in ways that builds confidence in decision-making contributes to better health literacy than education alone. The reasons mentioned above lead to the main objective of this research to develop health literacy of the older adult with the expectation that literacy will lead to appropriate decision-making and self-management. Health literacy is associated with self-management abilities, therefore, early recognition of low health literacy among older person together with interventions to improve health literacy might be very beneficial for older adults (Geboers, 2016).

Therefore, the objective of this research was to study the effectiveness of a health literacy program for self-management on food consumption among hypertensive older adult. The research framework employs the concept of health literacy (Nutbeam, 2000), namely access to information and knowledge, data analysis skills, utilizing information and knowledge, guiding of knowledge contributes to the health of others, and concept of self-management by assessing health needs, selecting use sources, and behavioural changes under health needs (Creer, 2000). There have been research supporting the study of the relationship between health literacy and self-management, health literacy and food consumption behavior (Geboers, 2016; Namdar et al., 2021). In this research, we explored these elements with the aim of viewing the effectiveness of the health literacy program.

## Materials and methods

### Study design

This quasi-experimental study was conducted using two groups and a pre-posttest design. The study period was from September 2019 to October 2020. Intervention conducted eight weeks and measured nutrition literacy, self-management on food intake, and food consumption pattern.

### Population and sample

The reference population was the older adult, both male and female, aged between 60 and 65 years, who lived in the Bang Phlat District, Bangkok, Thailand. We selected this particular age group for our implementation because this is an early retirement age group and chronic diseases start to affect their health. Hence, changing behaviour prior to serious health consequences is therefore an imperative measure for disease prevention.

Samples were drawn from the reference population, who met the inclusion criteria as follows: 1) between the ages of 60 - 65 years, male or female 2) have a blood pressure  $\leq 140/90$  mmHg 4) fluency in reading and writing Thai language 5) participate in the project voluntarily, and 6) have lived in the community for more than three months.

The sample size calculation used power analysis. A one-tailed test model was selected, and a method for comparing the mean scores used two independent group tests. The influence effect size was 0.80 (Meethien et al., 2011; Sawekwan, 2019), the confidence level was 0.05, and the test power was 0.95 (Cohen, 1992). The results of the calculation were a sample group of 70 people, 35 in the experimental group and 35 in the control group.

The sampling technique used a stratified multistage sampling method. Bang Phlat district consists of 4 sub-districts, namely Bang Phlat Sub-District, Bang Yi Khan Sub-District, Bang Ao Sub-District, and Bang Bamru District. Then communities from all four districts were selected to be represented at the provincial level using a probability proportional to

size. The communities that were in the area of the sample 4 communities were Daowadung Temple community (A), Bang Yi Khan Temple community (B), Chatkaew Jongkolnee Temple community (C), Khong Makham community (D). Community A and C were assigned experimental group using a non-random method. A total of 64 older adults met the inclusion criteria and were enrolled into our study. The number in the experimental was equal to the control group (32 in each group). Three people from each group withdrew during the course of the study. It is to be noted that sample of 32 from each group still provided 90% of study power.

#### Research instruments

Experimental research tool: Health literacy program. The activities design developed by researcher and used the concepts of health literacy of Nutbeam (2000), aimed to increase the subjects' health literacy and self-management and food consumption behavior. The details of the activities were making relationships, education, practicing communication skills by self-talk techniques, sharing experiences and learning, and evaluating the program by reflection from the group. The program conducted in eight weeks and it took 20 - 60 minutes to organize each activity (See Table 1). Kaewdamkerng (2019) stated that in general, habit is formed by completing a task continuing within 66 days (approximately 2 months) or 21 days at least. Moreover, previous research applied 6-8 weeks (Patipattarakul et al., 2018).

The program was conducted by researchers and held at each community. The program for experimental group comprised as following:

**Building relationship and health education:** Before we started health education, we built the relationship within the group to create trust, empowerment and encouragement within the group. These activities took 20 minutes. Also, during the week the health education was scheduled during the 1<sup>st</sup> week which lasted 60 minutes for small group teaching and discussion, via use of flip chart to demonstrate knowledge on hypertension, regarding: definition, causes, symptom, treatment, prevention and risk factors (WHO,2021; Thai Hypertension Society, 2019).

**Nutrition education:** The nutrition education was scheduled during 2<sup>nd</sup> week and lasted 60 minutes for small group teaching. We delivered the nutrition information related to healthy eating, a nutritional recommendation based on Dietary approaches to stop hypertension (DASH) (NIH, 2021), Thai Food Pyramid Guide, Dietary Guidelines, and Nutrition Facts Labels for Thai elders; essential nutrients (Nutrition Division, 2020) via picture media. We provided information on how to access resources and use of information (20 minutes for this session).

**Individual practice skill:** We demonstrate process of decision making by using tree diagram. Evaluation of decision-making was captured by asking questions about advantages or benefits, and disadvantages or risks of decisions. In addition, we demonstrated how to use such information, select resource, and verify the reliability of the information for example reading food label. After demonstration, they practiced their skill and showed researchers. We scheduled during the 3<sup>rd</sup> week of the program and took time 60 minutes of small group. The intervention materials included food label, leaflet, drug label etc.

**Practice self-communication skills:** Activities were scheduled during 4<sup>th</sup> week and it took 60 minutes. The activities consisted of overview of self-talk technique (Kross,2014; Health Direct, 2022) and practice self-talk by telling own messages to encourage and motivate subjects healthy eating and behaviour modification. Example of dialogues included: "I can do it I have accomplished something more difficult", "Can I eat salty food?" and "I have to read food labels before buying". The dialogues were developed by researchers based on review of existing information (Nutrition Division, 2020; Health Direct, 2022)

**Monitoring health eating:** Activities were scheduled during 5<sup>th</sup> and 6<sup>th</sup> weeks. Subjects were reported food consumption and self-talk technique to the researcher. It took approximately 20 minutes. We advised older adults whenever they were not appropriately performing healthy eating behavior.

**Sharing and Learning:** We delivered the activities during 7<sup>th</sup> week and took approximately 60 minutes of small group. Subjects were sharing experience and learning about self-talk technique, how to access health information and how to decision making and applied.

**Program evaluation:** We scheduled an evaluation by reflection during 8<sup>th</sup> week. This session took approximately 40 minutes for small group.

#### The intervention for control group

Like those in the experimental group, subjects in the control group received the same usual health care and routine health education activities from their respective community health care providers. However, subjects in the control group did not receive the 8 weeks health literacy program that was administered to those in the experimental group. Their usual health care and routine health education activities included; curative care, home health care and health disease prevention information.

**Table 1** health literacy program for self-management on food consumption

Session/Week	Activities	Relation between health literacy and self-management
<u>Week 1</u>		
Session 1: 20 mins	1. Building relationship within the group to create trust and empowerment and encouragement within the group	Functional health literacy (empowerment)
Session 2: 60 mins	1. Providing health education	
<u>Week 2</u>		
Session 3: 60 mins	1. Providing knowledge about nutrition for hypertension, DASH diet, diet control guidelines	Functional health literacy (access and understand)
Session 4: 20 mins	2. Knowledge about how to access resources and use of information	
<u>Week 3</u>		
Session 5: 60 mins	1. Training about how to decision making 2. Practice skills in selecting resources, and verifying the reliability of the information	Functional health literacy (access and understand)
<u>Week 4</u>		
Session 6: 60 mins	1. Overview and introduction self-talk 2. Practice self-communication skills by using the self-talk technique by telling own messages (internal dialogue) for behaviour modification	Interactive health literacy (analytic and problems solving) Self-management
<u>Week 5 – 6</u>		
Session 7: 20 mins by telephone	1. Individually reported on food consumption and self-talk technique to the researcher	Critical health literacy (decision making and apply) Self-management
<u>Week 7</u>		
Session 8: 60 mins	1. Experience sharing and learning (Sharing and Learning)	Functional health literacy (understand)
<u>Week 8</u>		
Session 9: 40 mins	1. Program evaluation	

The instrument for data collection comprised of four questionnaires and developed by literature review as follows:

- 1) The personal data questionnaire consists of seven items, namely sex, age, BMI, congenital disease, cigarette smoking, alcohol drinking, and physical activity.
- 2) The nutrition literacy questionnaire consists of 15 closed-ended questions. The answer is five-point on the Likert scale, i.e., never practice, seldom practice, sometimes practice, often practice, and always practice (75 full marks).
- 3) Food consumption patterns for the older adult questionnaire consists of 13 closed-ended questions. The answer is five-point on the Likert scale, i.e., never practice, seldom practice, sometimes practice, often practice, and always practice (65 full marks).
- 4) The questionnaire for self-management food intake pattern consists of 16 closed-ended questions. The answer is five-point on the Likert scale, i.e., never practice, seldom practice, sometimes practice, often practice, and always practice (80 full marks).

The determination of benchmarks to classify health literacy scores, dietary patterns, and self-management behaviours on food consumption used a 4-level classification based on the criteria for measuring the health quotient of the Health Education Division, Department of Health Service Support, Ministry of Public Health (2018): Poor <60% of the full score, Fair  $\geq 60\%$  - <70% of the full score, Good  $\geq 70\%$  - <80% of the full score, and Very Good  $\geq 80\%$  of the full score.

#### Measurement instrument

Health literacy program was validated by three experts in nutrition, adult and geriatric nursing, and community health nursing. By checking how well the results correspond to established theories and other measures of the same concept,

security and the possibility of using the program. Health literacy program was revised according to the recommendations before applying to the sample group. The Index of Item-Objective Congruence (IOC) was used so as to find the content validity. In this process, the questionnaires were checked by three experts including, one expert in nutrition, two experts in adult and geriatric nursing, and community health nursing field. The items that had scores lower than 0.5 were revised. On the other hand, the items that had scores higher than or equal to 0.5 were reserved. All of questionnaires has IOC between 0.8 to 1. The reliability of the questionnaires was determined to ensure that the responses collected through the instrument were reliable and consistent. The questionnaire was tested with 30 older persons that were not in the sample group. The reliability value was calculated by using Cronbach's alpha to ensure whether there was internal consistency within the items. According to the pre-test, the Cronbach's Alpha was 0.78, so the questionnaire was acceptable reliability (Tavakol & Dennick, 2011).

#### Ethical approval

This study was approved by the Ethical Review Committee for Human Research, Research and Development Institute, Suan Dusit University, Thailand (SDU-RDI 2020-008). Each patient who participated in the study was informed of the nature and objectives of the study. A written consent form was obtained before data collection from each participant.

#### Data Collection

Once the older adults, in both the experimental and controls groups, were identified, they were informed the purpose of the research. Appointments were made during the first week to collect data in their communities. Data were obtained from the older adults via administration questionnaires. We conducted the health literacy development program. The experimental group received the health literacy program, along with the usual health care and routine health education activities provided by their health care providers. The control group received only the usual health care and routine health education activities provided by their health care providers. During the 9<sup>th</sup> week, the questionnaires were administered to each older adults in each group. During the 10<sup>th</sup> week (9 weeks after the experimental group completed the health literacy program and all data were collected), the researchers provided the control group members the same health literacy program. Details of the protocol timeline is showed in Table 2.

**Table 2** Research protocol timeline

Week	0	1	2	3	4	5	6	7	8	9	10	
Experimental group												
measurement	T1	-	-	-	-	-	-	-	-	-	T2	
intervention	-	P <sub>1</sub>	P <sub>2</sub>	P <sub>3</sub>	P <sub>4</sub>	P <sub>5</sub>	P <sub>6</sub>	P <sub>7</sub>	P <sub>7</sub>	P <sub>8</sub>	P <sub>9</sub>	-
		Group					Individual		Group			
Control group												
measurement	T1	-	-	-	-	-	-	-	-	-	T2	HL
intervention	-	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	P <sub>0</sub>	

Note: 0 = Before beginning the experimental intervention  
 T1 = Pre-test; measurement of the nutritional literacy, food consumption, self-management before the experimental intervention  
 T2 = Post-test; measurement of the nutritional literacy, food consumption, self-management at 9<sup>th</sup> week  
 P<sub>0</sub> = Usual care  
 P<sub>1</sub> – P<sub>6</sub> = Group session 1-6  
 P<sub>7</sub> = Individual session  
 P<sub>8</sub> – P<sub>9</sub> = Group session 8-9  
 HL = Provision of the health literacy program

#### Data analysis

Descriptive statistics were used to summarize demographic data. Continuous variables were presented as mean ± standard deviation, and categorical variables were shown as frequencies and percentages. Chi-square, Fisher's exact test was used to compare the demographic data between group, and the independent t-test were used to examine differences between intervention and control groups, based on the assumptions of each statistic. Paired t-test was used to analyze in the difference between pre and posttest for the same subject. *P* values less than 0.05 were considered as statistical significance. Content analysis was used to summarize the results from the reflection.

## Results

A total, 64 participants completed the study, with 32 in the experimental and 32 in the control group. All characteristics of participants in both groups were similar in gender, age, cigarette smoking, alcohol drinking, and physical activity except BMI (Table 2).

The general characteristics of the experimental group was that most were female (84.4%). Participants' ages ranged from 60 to 65 years old, with a mean age of 63.34 years old (SD =1.89). Half the participants had BMI at the level of obesity (50.0 %). Twenty five percent of the participants had congenital diseases such as high blood pressure and diabetes. The majority (93.8%) were non-smokers and did not drink alcohol (84.4%). Only one person exercised daily (3.1%). The majority

of the control group was also females (68.8%). Their age ranged from 60 to 65 years old, with a mean age of 62.78 years old (SD = 1.93). The BMI at the level of obese was 43.8%. With regards to the control group, there were 37% of congenital diseases such as high blood pressure and diabetes. The majority of the participants were non-smokers and did not drink alcohol (84.4% and 65.6%, respectively). There was 9.4% exercise every day (Table 3). No significant differences were found between the demographic of the two group (p-value>0.05), except BMI and congenital diseases (p-value<0.05).

**Table 3** Demographic characteristics of participants

Characteristics	Experimental group (n = 32)		Control group (n = 32)		P-value
	n	%	n	%	
<b>Gender</b>					0.637
male	5	15.6	10	31.3	
female	27	84.4	22	68.8	
<b>Age (years)</b>					0.187
60 - 61	8	25.0	10	31.3	
62 - 63	5	15.7	7	21.9	
64 - 65	19	59.3	15	46.9	
	Mean =63.34 SD.=1.89		Mean =62.78 SD.=1.93		
<b>BMI (Kg./m.<sup>2</sup>)</b>					0.000
< 18.5	1	3.1	0		
18.5 - 22.9	8	25.0	6	18.8	
23.0 - 24.9	7	21.9	12	37.5	
≥ 25	16	50.0	14	43.8	
	Mean = 25.22 SD.=3.54		Mean = 24.87 SD.=2.62		
<b>Congenital diseases</b>					0.03
Yes	8	25.0	12	37.5	
No	24	75.0	20	62.5	
<b>Cigarette smoking</b>					1.00
Yes	1	3.1	1	3.1	
Recovering smoker	1	3.1	4	12.5	
No	30	93.8	27	84.4	
<b>Alcohol drinking</b>					0.762
Yes	1	3.1	1	3.1	
Recovering drinker	4	12.5	10	31.3	
No	27	84.4	21	65.6	
<b>Physical activity</b>					0.119
Everyday	1	3.1	3	9.4	
> 2 days - ≤ 5 days /week	15	46.9	12	37.5	
≤ 2 days /week	16	50.0	17	53.1	

The overall pre-test nutritional literacy of the experimental group and the control group was at a fair level meaning sufficient nutrition literacy and be able to practice it correctly. After the experiment, results found that most of the experimental groups had very good nutrition literacy and had proper and consistent practice. For the majority of the control group, there was a fair level of nutrition literacy.

The overall food consumption behaviour of the experimental group and the control group was at a fair level. In other words, participants had a minority of correct dietary consumption. After the experiment, results found that most of the experimental groups were at a good level of dietary behaviour. The majority of the control group had food consumption behaviour at a fair level.

The pre-test self-management behaviour on food consumption in the experimental group showed that most of subjects were at a fair level. They were able to manage themselves on food consumption well. The majority of the control group was poor. They could not manage their food consumption well. However, after the experiment, results suggested that most of the experimental group had a very good level of self-management behaviour. In the majority of control group, the level of self-management behaviour of food consumption was moderate. (Table 4)

**Table 4** Level of nutrition literacy, food consumption and self-management behaviours

	Scores	Level	Pre-test n (%)		Post -test n (%)		p-value
			Experimental group (n=32)	Control group (n=32)	Experimental group (n=32)	Control group (n=32)	
Nutrition literacy (75 full marks)	< 60% of full marks	Poor	10 (31.3)	12 (37.5)	0	11 (34.4)	0.000
	≥ 60% - <70% of full marks	Fair	12 (37.5)	17 (53.1)	2 (6.3)	19 (59.4)	
	≥ 70% - <80% of full marks	Good	10 (31.3)	3 (9.4)	11 (34.4)	2 (6.3)	
	≥ 80% of full marks	Very good	0	0	19 (59.4)	0	
Food consumption behaviour (65 full marks)	< 60% of full marks	Poor	11 (31.4)	7 (21.9)	2 (6.3)	7 (21.9)	0.346
	≥ 60% - <70% of full marks	Fair	14 (43.8)	20 (62.5)	8 (25.0)	20 (62.5)	
	≥ 70% - <80% of full marks	Good	7 (21.9)	5 (15.6)	13 (40.6)	5 (15.6)	
	≥ 80% of full marks	Very good	0	0	9(28.1)	0	
Self-management behaviours (80 full marks)	< 60% of full marks	Poor	7 (21.9)	18 (56.3)	0	6 (18.8)	0.451
	≥ 60% - <70% of full marks	Fair	17 (53.1)	13 (40.6)	1 (3.1)	19 (59.4)	
	≥ 70% - <80% of full marks	Good	8 (25.0)	1 (3.1)	6 (18.8)	7 (21.9)	
	≥ 80% of full marks	Very good	0	0	25 (78.1)	0	

The experimental group had the mean score of all components (nutritional literacy, food consumption and self-management) significantly higher than the control group with statistical significance at level of <.001. (Table 5).

**Table 5** the mean score of nutritional literacy, food consumption behavior, and self-management behavior between experimental and controlled group by using independent t-test

variables	Experimental group		Controlled group		p-value
	mean	SD.	mean	SD.12	
<b>Before</b>					
nutritional literacy	47.72	6.0	45.31	5.29	0.940
food consumption behavior	40.06	4.85	40.40	3.32	0.742
self-management behavior	51.91	6.26	66.66	5.77	0.001
<b>After</b>					
nutritional literacy	60.25	5.63	45.38	5.19	<0.001
food consumption behavior	47.53	5.58	40.81	3.27	<0.001
self-management behavior	66.65	5.77	52.06	5.69	<0.001

The experimental group had the mean score of all components significantly higher than before experiment with statistical significance at level of  $<.001$ . (Table 6).

**Table 6** the mean score of nutritional literacy, food consumption behavior, and self-management behavior between before and after intervention of experimental group by using dependent t-test

variables	Before experiment		After experiment		p-value
	mean	SD.	mean	SD.12	
nutritional literacy	47.72	6.0	60.25	5.63	$<0.001$
food consumption behavior	40.06	4.85	47.53	5.58	$<0.001$
self-management behavior	51.91	6.26	66.65	5.77	$<0.001$

#### Evaluation of the program by reflection from the experimental group

The results from the reflection of the experimental group were summarized as follows: the understanding of the information or health literacy for self-management on food consumption must be studied and understood well; read more; do not believe in delivered information without factual information from knowledgeable people; or from advertising with solicitation to buy. People and communities with access to information and health literacy need to seek further knowledge and build empowerment, community and social support, and effective communication at the family level, as well as expand widely at the community level. People must interpret information before they can analyze and make food consumption decisions. Therefore, having academics come to suggest and help develop community leadership skills results in an analytical thinking process to be a leader in community development and improve people in the community to gain skills. The methods of developing health literacy for self-management on food consumption were health education from academics giving direct knowledge. People can transmit the knowledge correctly to the community. In addition, there are self-talk techniques, communication in small groups, and communication at the community level. Self-talk is a method that helps people be mindful before doing anything. It serves as a reminder to them every time they do something, until they become second nature to them. For example, they read a nutrition label before buying a product or searching for information. Moreover, sharing experience and learning support health literacy and self-management, such as exchanging knowledge on how to self-manage and make decisions on consumption. The visual teaching materials are important to enhance health literacy for self-management in groups with learning restrictions. Therefore, the use of illustrations and explanations is suitable for the older adult, such that pictures can provide more understanding.

#### Discussion

All characteristics of participants in both groups were similar in gender, age, cigarette smoking, alcohol drinking, and physical activity. However, BMI showed significant difference at level of  $<0.001$ , and the mean score of BMI experimental group greater than control group. The finding suggested that the experimental group showed a great self-management. This indicated that BMI factor could have an effect on self-management. Vinkers et al. (2014) suggested that good self-management intervention in overweight could improve outcomes in behavioral and anthropometric. This indicated that older adult's obesity might have greater interest in program to improve their health.

The level of nutrition literacy and self-management behaviour in the experimental group increased after the experiment. It has been suggested that the development of health literacy, guidelines, and appropriate activities can develop and promote health literacy (WHO, 2009). Good health literacy will improve health behaviours (Kingkaew and Prasertsri, 2015). Likewise, Rattanawarang and Chanta (2018) reported a study that health literacy is related to self-care behaviour in chronic disease, and the testing program showed the mean scores on nutrition literacy increased significantly after testing of the experimental group ( $p\text{-value} < 0.001$ ). Our study findings were supported by several previous studies (Thep-in, 2019; Boontanon et al., 2019; Visscher et al., 2018). An appropriate activity for the group will improve health literacy (WHO, 2009). Methods and guidelines for development in organizing the learning process led to learner memorization learning (Stacey et al., 2011; Visscher et al., 2018; Kaewdamkeeng, 2019). The results of the mean score before and after the experiment of food consumption behaviour and self-management behaviour on food intake of the experimental group showed a significantly higher score than before the test ( $p\text{-value} < 0.001$ ). The results could be explained by knowledgeable individuals, having a variety of information which helps them making decisions about behaviour improvement (Pongkiatchai & Wongwisesskul, 2018). Moreover, Santo et al. (2005) stated that the development of health literacy, especially health education following by group or individual practices, will help promote decision-making skills on health behaviours and good self-management. Many studies have found that health literacy is associated with positive self-management and health behaviours (Kingkaew & Prasertsri, 2015; Wang et al., 2017; Rattanawarang & Chantha, 2018). The study by Jayasinghe (2016) found that low health literacy had a significant effect on health behaviours among Australian patients ( $p\text{-value} < 0.001$ ). Therefore, in our study the effects of the health literacy development program on the older adults for self-management in food consumption made the experimental group more nutrition literacy, consumption behaviour and self-management behaviour. The experimental group's mean scores were significantly different from the control group's ( $p\text{-value} < 0.001$ ). Our findings suggested that health literacy and health education, including activities to promote knowledge, can empower individuals to self-management in terms of health and the ability to make decisions about their health more appropriately.

Interestingly, the evaluation of reflections toward activity among experimental group indicated that perspectives of older adults on nutritional health literacy appeared to be associated with understanding of the information, information accessibility, the ability to transform the information to daily life practice. Likewise, older adults reflected the emphasized



strategies to augment health literacy helped them with self-management on nutritional habits to improve their health. These strategies included education, communication, sharing experiences, and infographic or illustration. Moreover, the ability to read and write are important for older adults to gain and understand the information on basic health literacy, especially in health education and cognitive appraisal on health information such as the predominated risk, medical information, and nutrition labels (Nutbeam, 2000; Sorensen, et al., 2012; Pongkiatchai & Wongwisukul, 2018). Health information is typically available and older adult could seek the information resources. However, the complexities of the different data structures influence on the decision to evaluate and apply information (Harzheim et al., 2020). World Health Organization (WHO) implied that readability, transformation skills, and an insightful understanding are the principal factors for enhancing health literacy and improving an effective communication among individuals (WHO, 2009). Besides, information accessibility is an interactive health literacy (Nutbeam., 2000; Sorensen et al., 2012; Pongkiatchai & Wongwisukul, 2018). It is ability to find a strategy to promote self-management by seeking and classifying information across the different sources. Ghaffari-Fam et al. (2020) studied health information and blood pressure control. They found that the accessibility to seek health information could predict blood pressure control among Iranian hypertensive clients (p-value < 0.05). In sum, the availability, accessibility of the information and communication are crucial to generate nutritional health literacy.

The ability to apply knowledge skills are a critical health literacy level (Nutbeam, 2000; Sorensen et al., 2012; Pongkiatchai & Wongwisukul, 2018). The older adult ability to interpret data, analyzing and decision-making lead to make decisions in self-management. The results of the activity assessment showed that knowledge and understanding of the older adult helped making a decision and ability to apply information. This study findings are in keeping with the results of Boontanon et al., (2019). They reported that the older adult with good health literacy led to nutrition decision making and behavioural change. Besides, a study by Ghaffari-Fam et al. (2020) found that people with healthy literacy were able to control high blood pressure.

When discussing about the promoting health literacy for self-management in food consumption, the older adult pointed that educating, communicating, sharing experiences and using infographic illustration could improve health literacy. This suggestion is in line with the Kwanmuang's study (Kaewdamkereng, 2019) which reported that the strategies of promoting health literacy are varied based on the health literacy level such as the functional health literacy level, the strategies of promoting health literacy, the learning process development and the educational tool selection. Moreover, the interactive health literacy level, such as the questioning skills and creating a friendly atmosphere for learning, and the critical health literacy level, such as the method of promoting health literacy, the training in decision-making skills, identifying problems, searching for information to help evaluate and decide, self-assessment and self-management are also included. According to the health education study results from Santo et al (2005), it was mentioned that the development of health literacy, particularly health promotion as well as individual and group practice will enhance the decision-making skills toward health behaviours and self-management. Allen et al (2017) stated that health lifestyle will be improved by the individual or group health promotion as well as launching group activities, sharing experiences and discussing topics related to communication channel that the information is reliable and of high-quality. The communication in this research is intrapersonal communication or self-talk, following mindfulness in Buddhist psychology way which was used amongst the athletes. Tod et al. (2011) suggested that positive self-talk can motivate self-empowerment. It also helps the athletes to be more concentrated during competitions. The findings were supported by the psychologist as self-talk helps people to consolidate positive thoughts, feelings, emotions in stressful situations. (Kross et al., 2014). It is similar to the Buddhism (constant mindfulness) as it induces precepts, meditation and reduce stress. For the instructional tools, our results are consistent with the study of Kaewdamkereng (2019) in that the process of promoting health literacy, is essential to choose appropriate and standard instructional medias. To our knowledge, there are several instructional tools available, however these are used in many ways based on the type of study objectives and participants. In this current study, we employed image and explanation for the older adult, as this type of tool allows the older adult to ask questions and also helps the researcher to clarify any points that the older adult was not fully understood. There was an agreement between researchers and participants on using images as a teaching media as well.

This study has some limitations including measured food consumption behaviour and self-management behavior for only at a short period. Secondly, this study did not follow blood pressure control, which is an important clinical outcome. Therefore, it could not be assured whether this program was effective in blood pressure control. The potential effectiveness of health literacy program and health outcomes would be a relevant topic for future research with long period study.

## Conclusion

In conclusion, a health literacy development program for self-management on food consumption of the older adult can increase the level of nutrition literacy, dietary behavior, and self-management. Besides, the method to enhance health literacy for self-management of food consumption is educating, communicating, sharing experiences, and use the Fotonovela technique. It could develop health literacy in the older adult as well. Therefore, we recommended that it would be suitable to implement at primary public health center to improve health literacy among older person. They could also use self-talk technique and sharing learning to improve decision making and applied skills.

## Acknowledgement

We extend our appreciation to all study participants. This work was supported by grant from The Suan Dusit University.

## References

Allen, M.P., Auld, E., Logan, R., Montes, J.H. (2017). Improving collaboration among health communication, health education, and health literacy. Retrieved from <https://www.rootcausecoalition.org/wp->

- content/uploads/2017/08/Improving-Collaboration-among-Health-Communication-Health-Education-and-Health-Literacy.pdf
- Bellisle, F. (2019). The value of studying laboratory meals. The effects of environment on product design and evaluation pages 209-225. Retrieved from <https://doi.org/10.1016/B978-0-12-814495-4.00010-6>
- Boontanon, N., Armok, S., Surapattanachai, N. (2019). The effectiveness of nutrition literacy promoting for the older adult in outpatient unit, Health Promotion Hospital regional health promotion center 9, Nakhon Ratchasima. *Journal of Health Promotion Center 9: Journal of Health Promotion and Environmental Health*, 13(30), 1-14.
- Chodosh, J., Morton, S.C., Mojica, W., Maglione, M., Suttrop, M.J., Hilton, L., Rhodes, S., Shekelle, P. (2005). Meta-analysis: Chronic disease self-management programme for older adults. *Annual of Internal Medicine*, 143(6), 427-439. Chao, et al., 2013;
- Creer, L.T. (2000). Self-management of chronic illness. *Handbook of self-regulation*. California: Academic. 601-629.
- Division of non-communicable diseases. (2022). Mortality with 5 NCDs. Retrieved from <http://www.thaincd.com/2016/mission/documents-detail.php?id=14220&tid=32&gid=1-020>
- Ekplakorn W., (editor). (2014). *Thai National Health Examination Survey (NHES V)*. Nonthaburi: The Graphico system co.Ltd.
- Geboers, B., de Winter, A. F., Spooenberg, S. L., Wynia, K., & Reijneveld, S. A. (2016). The association between health literacy and self-management abilities in adults aged 75 and older, and its moderators. *Quality of life research: an international journal of quality-of-life aspects of treatment, care and rehabilitation*, 25(11), 2869–2877. Retrieved from <https://doi.org/10.1007/s11136-016-1298-2>
- Ghaffari-Fam, S., Sarbazi, E., Ardabili, F., Babazadeh, T., Darghahi, G., Jafaralilou, H. (2020). The role of health literacy in hypertension control: a cross-sectional study in Iran. *Ann Ig*, 32(3), 263-273.
- Harzheim, L., Lorke, M., Wooten, C., Jünger, S. (2020). Health Literacy as Communicative Action—A qualitative study among persons at risk in the context of predictive and preventive medicine. *Int J Environ Res Public Health*, 17(5), 1718.
- Health Direct. (2022). Self-talk. Retrieved from <https://www.healthdirect.gov.au/self-talk#:~:text=Self%20talk%20is%20the%20way,internal%20monologue%20throughout%20the%20day>.
- Health promotion glossary. (1998). World Health organization. Retrieved from [www.who.int/healthpromotion/about/HPR%20Glossary%201998.pdf](http://www.who.int/healthpromotion/about/HPR%20Glossary%201998.pdf).
- Jayasinghe, U.W., Harris, M.F., Parker, S.M. et al. (2016). The impact of health literacy and life style risk factors on health-related quality of life of Australian patients. *Health Qual Life Outcomes*, 14, 68. Retrieved from <https://doi.org/10.1186/s12955-016-0471-1>
- Kaewdamkerng, K. (2018). *Health literacy: access, understanding and application*. Second printing. Bangkok: Amarin Printing and Publishing, Company Limited.
- Kaewdamkerng, K. (2562). 21 days really change behavior. Retrieved from <https://shorturl.asia/k02SU>
- Kim, S., Love, F., Quistberg, D.A., Shea, J.A. (2004). Association of health literacy with self-management behavior in patients with diabetes. *Diabetes Care*, 27(12), 2980-2983.
- Kross, E., Bruehlman-Senecal, E., Park, J., Burson, A., Dougherty, A., Shablack, H., Bremner, R., Moser, J., & Ayduk, O. (2014). Self-talk as a regulatory mechanism: How you do it matters. *Journal of Personality and Social Psychology*, 106(2), 304–324. Retrieved from <https://doi.org/10.1037/a0035173>
- Meethien, N., Pothiban, L., Ostwald, S.K., Sucamvang, K., Panuthai, S. (2011). Effectiveness of nutritional education in promoting healthy eating among elders in Northeastern Thai. *Pacific Rim Int J Nurs Res*, 15(3), 188-202.
- Mohd Isa, D.; Shahar, S.; He, F.J.; Majid, H.A. Associations of health literacy with blood pressure and dietary salt intake among adults: A systematic review. *Nutrients* 2021, 13, 4534. Retrieved from <https://doi.org/10.3390/nu13124534>
- National Committee on the Older adult, Department of Older adults, Ministry of Social Development and Human Security. (2020). *The Action Plan for the Older adult, Phase 2 (2002 - 2022)*, revised 2<sup>nd</sup> . Retrieved from [https://www.dop.go.th/download/laws/th1614675775-828\\_0.pdf](https://www.dop.go.th/download/laws/th1614675775-828_0.pdf).
- NIH (National Heart, Lung and Blood institute). (2021). DASH eating plan. Retrieved from <https://www.nhlbi.nih.gov/education/dash-eating-plan>
- Norris S.L., Lau J., Smith J., Schmid C.H., Engelgau, M.M. (2002). Self-management education for adults with type 2 diabetes: a meta-analysis of the effect on glycemic control. *Diabetes Care*, 25(7), 1159-1171.
- Nutbeam, D. (2000). Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health Promotion International*, 15(3), 259-267.
- Nutbeam, D. (2010). The evolving concept of health literacy. Retrieved from <http://www.bumc.bu.edu/healthliteracyconference/files/2010/04/Don-Nutbeam.pdf>. Sangduean Kingkaew and Nusara Prasertsri, (2015).
- Nutrition Division, Ministry of Public Health of Thailand. (2022). Food and exercise for older persons: Nutritional guideline and nutritional flag. Retrieved from <https://nutrition2.anamai.moph.go.th/th/book/194497>
- Patipattarakul, P., Korjedee, P., Pongsakchart, P., Yingrengreung, S., Burapa, C., Jakpon, P. (2018). The effectiveness of a health promotion program on health behaviors of elderly in community. *Journal of Health Science Research*, 12 (supp), 52-60.
- Pongkiatchai, R. & Wongwisetkul, S. (2018). Nutrition literacy and the older adult with hypertension. *Food Health and Bioenvironmental Science*, 11(3), 49-55.
- Rattanawarang, W. & Chantha, W. (2018). Health literacy of self-care behaviors for blood glucose control in patients with type 2 diabetes, Chainat Province. *The Journal of Baromarajonani College of Nursing, Nakhonratchasima*, 24(2), 34-51.

- Santo, A., Laizner, A.M., & Shohet, L. (2005). Exploring the value of audiotapes for health literacy: a systematic review. *Patient Education and Counseling*, 58(3), 235-243
- Sawekwan, J., Amnatsatsue, K., Kerdmongko, P. (2019). Effects of a continuing care program for older adults with uncontrolled hypertension. *Journal of Public Health Nursing*, 33(1), 152-170.
- Sorensen, K., Van den Broucke, S., Fullam, J., Doyle†, G., Pelikan, J., Slonska, Z., & Brand, H. (2012). Health literacy and public health: A systematic review and integration of definitions and models. *BMC Public Health*, 12(80), 1-13.
- Stacey, L., Sheridan, D.J., Halpern, A.J., Viera, N.D., Berkman, K.E., Donahue & Karen Crotty. (2011). Interventions for individuals with low health literacy: A systematic review, *Journal of Health Communication*, 16(sup3), 30-54, DOI: 10.1080/10810730.2011.604391
- Steverink N, Lindenberg S, Slaets JPJ. (2005). How to understand and improve older people's self-management of wellbeing. *European Journal of Ageing*, 2(4), 235–244. Retrieved from [https://doi: 10.1007/s10433-005-0012-y](https://doi.org/10.1007/s10433-005-0012-y)
- Tavakol, M., & Dennick, R. (2011). Making sense of Cronbach's alpha. *International journal of medical education*, 2, 53–55. Retrieved from <https://doi.org/10.5116/ijme.4dfb.8dfd>
- Thai Hypertension Society. (2019). Thai Guidelines on The Treatment of Hypertension. Retrieved from <http://www.thaihypertension.org/files/HT%20guideline%202019.with%20watermark.pdf>
- Thepin, K., Moolsart, S., Jantacumma, N. (2019). Effectiveness of a Health Literacy Developmental Program in Patients with Uncontrolled Hypertension. *Boromarajonani College of Nursing, Uttaradit Journal*, 11(1), 197-212.
- Tod, D., Hardy, J., Oliver, E. (2011). Effects of self-talk: A systematic review. *Journal of Sport & Exercise Psychology*, 33, 666-687.
- Van de Velde, D., De Zutter, F., Satink, T., Costa, U., Janquart, S., Senn, D., & De Vriendt, P. (2019). Delineating the concept of self-management in chronic conditions: a concept analysis. *BMJ open*, 9(7), e027775. <https://doi.org/10.1136/bmjopen-2018-027775>
- Velardo S. (2015). The nuances of health literacy, nutrition literacy, and food literacy. *J Nutr Educ Behav*, 47(4), 385–9. Retrieved from <https://doi.org/10.1016/j.jneb.2015.04.328>.
- Vinkers, C.D.W., Adriaanse, M.A., Kroese, F.M. *et al.* (2014). Efficacy of a self-management intervention for weight control in overweight and obese adults: a randomized controlled trial. *J Behav Med* 37, 781–792. Retrieved from <https://doi.org/10.1007/s10865-013-9530-9>
- Visscher, B.B., Steunenberg, B., Heijmans, M., Hofstede, J.M., Devillé, W., Iris van der Heide, Rademakers J. (2018). Evidence on the effectiveness of health literacy interventions in the EU: a systematic review. *BMC Public Health*, 18, 1414.
- Wang, C., Lang, J., Xuan, L., Li, X., & Zhang, L. (2017). The effect of health literacy and self-management efficacy on the health-related quality of life of hypertensive patients in a western rural area of China: a cross-sectional study. *International journal for equity in health*, 16(1), 58. Retrieved from <https://doi.org/10.1186/s12939-017-0551-9>
- WHO. (2009). Promoting Health and Development: Closing the Implementation Gap, Nairobi, Kenya. Retrieved from [https://www.who.int/healthpromotion/conferences/7gchp/Track1\\_Inner.pdf](https://www.who.int/healthpromotion/conferences/7gchp/Track1_Inner.pdf).
- WHO. (2021). Hypertension. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/hypertension>
- World Health Organization. (2013). A global brief on hypertension: Silent killer, global public health crisis: World Health Day. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/79059/W?sequence=1>
- Wongwiseskul, S. & Chinnapan, W. (2014). Program of health education through group learning and 'buddy' support on reducing excessive salt consumption among the aging with hypertension. Proceedings of the 1st International Nursing Conference (INC 2014) Nursing Innovation: A key to promote the humanized health in ASEAN Economic Community (AEC) Era. Ho Chi Minh City, Vietnam.